

Modular Constraints on Conformal Field Theories with Currents

Jin-Beom Bae

Korea Institute for Advanced Study

Based on ArXiv:1708.08815
work with Sungjay Lee and Jaewon Song

Autumn Symposium on String Theory
2017. 9. 11

Introduction

Conformal Field Theories in Higher Dimension ($d > 2$)
Representation : $SO(d, 2)$ with generator $P_\mu, K_\mu, M_{\mu\nu}, D$
 Δ (continuous), ℓ (discrete)

Conformal Field Theories in Two dimension
Governed by the Virasoro generator L_n and \bar{L}_n , $n \in \mathbb{Z}$
 h, \bar{h} ($\Delta = h + \bar{h}$ and $\ell = |h - \bar{h}|$)

$c < 1$

$c > 1$

- Unitary rep defined only for

$$c = 1 - \frac{6}{m(m+1)},$$

$$h_{r,s} = \frac{[(m+1)r - ms]^2 - 1}{4m(m+1)}$$

- Finite number of primaries :

Rational Conformal Field Theory

- Completely solved (integrable)

- Unitary rep defined for arbitrary h and \bar{h} (Similar to $d > 2$ CFTs)

- RCFTs (e.g., WZW models)

- Also includes **Irrational CFTs** (e.g., Liouville theories)

- A full classification of such theories still out of reach

Settings and Numerical Results

- The Character Decomposition

- The (Virasoro) vacuum characters and primary characters are defined by

$$\chi_0(\tau) = \frac{1}{\eta(\tau)} q^{-\frac{c-1}{24}} (1 - q), \quad \chi_h(\tau) = \frac{1}{\eta(\tau)} q^{h - \frac{c-1}{24}}$$

The torus partition function of unitary CFT admit the **character decomposition**,

$$Z(\tau, \bar{\tau}) = \chi_0(\tau) \bar{\chi}_0(\bar{\tau}) + \sum_{h, \bar{h}} d(h, \bar{h}) \chi_h(\tau) \bar{\chi}_{\bar{h}}(\bar{\tau}) + \sum_{j=1} \left[d(j) \chi_j(\tau) \bar{\chi}_0(\bar{\tau}) + \tilde{d}(j) \chi_0(\tau) \bar{\chi}_j(\bar{\tau}) \right],$$

where the degeneracies $d(h, \bar{h})$, $d(j)$ and $\tilde{d}(j)$ are positive integers.

- The constraints from $SL(2, \mathbb{Z})$

- **T- transformation** : All states should have **integer spin**.
- **S- transformation** : $Z(\tau, \bar{\tau}) = Z(-\frac{1}{\tau}, -\frac{1}{\bar{\tau}})$

$$\mathcal{Z}_0(\tau, \bar{\tau}) + \sum_{h, \bar{h}} d(h, \bar{h}) \mathcal{Z}_{h, \bar{h}}(\tau, \bar{\tau}) + \sum_{j=1} \left[d(j) \mathcal{Z}_j(\tau, \bar{\tau}) + \tilde{d}(j) \mathcal{Z}_{\tilde{j}}(\tau, \bar{\tau}) \right] = 0$$

where the function $\mathcal{Z}_\lambda(\tau, \bar{\tau})$ is defined as $\chi_\lambda(\tau) \bar{\chi}_\lambda(\bar{\tau}) - \chi_\lambda(-\frac{1}{\tau}) \bar{\chi}_\lambda(-\frac{1}{\bar{\tau}})$.

- Modular Bootstrap - Basic Strategy [Rattazzi, Rychkov, Tonni, Vichi 08], [Poland, Simmons-Duffin 10]

- In the computation, we mainly use the **reduced character** for convenience.

$$\hat{\chi}_0(\tau) = \tau^{\frac{1}{4}} \eta(\tau) \chi_0(\tau), \quad \hat{\chi}_h(\tau) = \tau^{\frac{1}{4}} \eta(\tau) \chi_h(\tau)$$

- Apply the linear functional $\alpha \left[\hat{\mathcal{Z}}(z, \bar{z}) \right] \equiv \sum_{m,n}^{m+n=N} \alpha_{m,n} \partial_z^m \partial_{\bar{z}}^n \hat{\mathcal{Z}}(z, \bar{z})$ to the modular bootstrap equation. ($\tau \equiv ie^z$, the crossing point at $z = 0$)

$$\alpha \left[\hat{\mathcal{Z}}_0(z, \bar{z}) \right] + \sum_{j=1}^{j_{\max}} \left(d(j) \alpha \left[\hat{\mathcal{Z}}^j(z, \bar{z}) \right] + \bar{d}(j) \alpha \left[\hat{\mathcal{Z}}^{\bar{j}}(z, \bar{z}) \right] \right) + \sum_{h, \bar{h} \in \mathcal{P}} d(h, \bar{h}) \alpha \left[\hat{\mathcal{Z}}^{h, \bar{h}}(z, \bar{z}) \right] = 0.$$

- Find $\alpha_{m,n}$ such that,

$$\alpha \left[\hat{\mathcal{Z}}_0(z, \bar{z}) \right] > 0,$$

$$\text{and } \alpha \left[\hat{\mathcal{Z}}^j(z, \bar{z}) \right] \geq 0, \quad \alpha \left[\hat{\mathcal{Z}}^{\bar{j}}(z, \bar{z}) \right] \geq 0 \quad \text{for } j \in \mathbb{Z},$$

$$\text{and } \alpha \left[\hat{\mathcal{Z}}^{h, \bar{h}}(z, \bar{z}) \right] \geq 0 \quad \text{for } (h, \bar{h}) \in \mathcal{P}$$

If we find such $\alpha_{m,n}$, then **we conclude that no modular invariant partition function can exist**. This problem can be converted to the **semi-definite programming**.

- Assumptions on the spectrum [Collier, Lin, Yin 16]

- In the modular bootstrap equation, we sum the primaries $(h, \bar{h}) \in \mathcal{P}$. We can make three different assumptions on \mathcal{P} .

Scalar Gap Problem

In this problem, we impose a gap Δ_s only to the scalar operator.

$$\begin{aligned} \Delta &\geq \Delta_s \text{ for } j = 0, \\ \Delta &\geq j \text{ for } j \neq 0. \end{aligned}$$

Overall Gap Problem

In this problem, we impose a gap Δ_o to the certain low-spin operators.

$$\Delta \geq \text{Max}(j, \Delta_o)$$


Twist Gap Problem

In this problem, we impose a gap Δ_t to the twist, defined as


$$\begin{aligned} t &\equiv \Delta - j. \\ \Delta &\geq j + \Delta_t \end{aligned}$$

- Additionally, **we impose the contribution of conserved currents** in the modular bootstrap equation.
- For a given c and Δ_{gap} (Δ_s or Δ_o or Δ_t), examine if one can find the numerical solution $(\alpha_{m,n})$ to the semi-definite programming or not. The results of this scanning process can be summarized on the two-dimensional plot.

• The Numerical Results ($c \leq 8$)



- The Numerical Results ($c \leq 8$), Focus on the Twist Gap



- Expected CFTs on the numerical bound (Twist Gap)
 - For the Wess-Zumino-Witten model with affine Lie algebra $\hat{\mathfrak{g}}$ and level- k ,


$$c = \frac{k \dim \hat{\mathfrak{g}}}{k + h^\vee}, \quad h_\lambda = \frac{(\lambda, \lambda + 2\rho)}{2(k + h^\vee)}$$

- The above formulae suggest that the **twist gap problem realize level-1 WZW models on the numerical boundary!** ($c \leq 8$)

Central Charge	Lowest Primary	Expected CFT
$c = 1$	$\Delta_t = 1/2$	$SU(2)_1$ WZW model
$c = 2$	$\Delta_t = 2/3$	$SU(3)_1$ WZW model
$c = 14/5$	$\Delta_t = 4/5$	$(G_2)_1$ WZW model
$c = 4$	$\Delta_t = 1$	$SO(8)_1$ WZW model
$c = 26/5$	$\Delta_t = 6/5$	$(F_4)_1$ WZW model
$c = 6$	$\Delta_t = 4/3$	$(E_6)_1$ WZW model
$c = 7$	$\Delta_t = 3/2$	$(E_7)_1$ WZW model
$c = 8$	$\Delta_t = 2$	$(E_8)_1$ WZW model

- The eight simple Lie group $A_1, A_2, G_2, D_4, F_4, E_6, E_7$ and E_8 are referred to as **Deligne's exceptional series**.

- The Numerical Result (Twist Gap, $c \leq 54$)



- The Numerical Result (Twist Gap, $c \leq 54$)
 - When the holomorphic currents are included from $j = 1$, the following four classes are further realized on the numerical boundary.

Central Charge	Lowest Primary	Expected CFT
$c = 16$	$\Delta_t = 2$	$(E_8 \times E_8)_1$ WZW model

- When the holomorphic currents are included from $j = 2$,

Central Charge	Lowest Primary	Expected CFT
$c = 24$	$\Delta_t = 4$	Monster CFT
$c = 48$	$\Delta_t = 6$	" $c = 48$ ECFT"
$c = 8$	$\Delta_t = 1$	CFT with $O_{10}^+(2)$
$c = 16$	$\Delta_t = 2$	CFT with $O_{10}^+(2)$
$c = 47/2$	$\Delta_t = 3$	Baby Monster CFT

- For instance, the unique modular invariant partition function at $c = 24$ is,

$$\begin{aligned}
 Z_{k=1}(q, \bar{q}) &= (j(q) - 744)(\bar{j}(\bar{q}) - 744) \\
 &= (1 + 196884q^2 + \cdots)(1 + 196884\bar{q}^2 + \cdots)
 \end{aligned}$$

- The Modular Differential Equation(MDE)

- Idea : n characters of rational conformal field theory(RCFT) are the solutions to the n -th order modular differential equation, [Mathur, Mukhi, Sen 88]

$$D_{\tau}^n \chi(\tau) + \sum_{k=0}^{n-1} \phi_k(\tau) D_{\tau}^k \chi(\tau) = 0,$$

with $D_{\tau} f(\tau) \equiv \partial_{\tau} f(\tau) - \frac{\pi i r}{6} f(\tau)$. (r is the modular weight of the test function $f(\tau)$)

- Second Order Modular Differential Equation

- To get the vacuum character, solve the second order differential equation,

$$D_{\tau}^2 \chi(\tau) + \hat{\mu} E_4(\tau) \chi(\tau) = 0,$$

with an ansatz $\chi_{\hat{\lambda}}(q) = q^{\alpha} (a_0 + a_1 q + a_2 q^2 + a_3 q^3 + a_4 q^4 + \dots)$.

- The coefficients $\{a_0, a_1, a_2, \dots\}$ are **positive integer** only for [Mathur, Mukhi, Sen 88], [Tuite 08]

$$c \in \left\{ \frac{2}{5}, 1, 2, \frac{14}{5}, 4, \frac{26}{5}, 6, 7, \frac{38}{5}, 8 \right\}.$$

- Third Order Modular Differential Equation

- To get the vacuum character, solve the third order differential equation,

$$D_\tau^3 \chi(\tau) + \mu_1 E_4(\tau) D_\tau \chi(\tau) + \mu_2 E_6(\tau) \chi(\tau) = 0,$$

with an ansatz $\chi_\lambda(q) = q^\alpha (a_0 + a_2 q^2 + a_3 q^3 + a_4 q^4 + \dots)$.

- The coefficients $\{a_0, a_2, a_3, \dots\}$ are **positive integer** only for [Mathur, Mukhi, Sen 88], [Tuite 08]

$$c \in \left\{ -\frac{44}{5}, 8, 16, \frac{47}{2}, 24, 32, \frac{164}{5}, \frac{236}{7}, 40 \right\}.$$

- The primary characters have the form of

$$\chi_{h_\pm}(\tau) = q^{h_\pm - \frac{c}{24}} \left[b_0 + b_1 q + b_2 q^2 + \dots \right]$$

with $h_\pm(c) = \frac{c+4}{16} \pm \frac{\sqrt{368+24c-c^2}}{16\sqrt{31}}$.

- The coefficients in the primary characters are **not completely fixed** from the modular differential equation.

Spectral Analysis

- Finding the degeneracy bound [Rattazzi, Rychkov, Vichi 10]

- Rewrite the modular bootstrap equation as

$$\alpha \left[\hat{\mathcal{Z}}_0(z, \bar{z}) \right] + d(h^*, \bar{h}^*) \alpha \left[\hat{\mathcal{Z}}^{h^*, \bar{h}^*}(z, \bar{z}) \right] + \alpha \left[\hat{\mathcal{Z}}^{rest}(z, \bar{z}) \right] = 0,$$

$$\alpha \left[\hat{\mathcal{Z}}^{rest}(z, \bar{z}) \right] \equiv \sum_{j=j_{min}}^{j_{max}} \left(d(j) \alpha \left[\hat{\mathcal{Z}}^j(z, \bar{z}) \right] + \bar{d}(j) \alpha \left[\hat{\mathcal{Z}}^{\bar{j}}(z, \bar{z}) \right] \right) + \sum_{h, \bar{h} \in \mathcal{P}} d(h, \bar{h}) \alpha \left[\hat{\mathcal{Z}}^{h, \bar{h}}(z, \bar{z}) \right],$$

and solve the following problem via the semi-definite programming.

$$\text{Maximize } \alpha \left[\hat{\mathcal{Z}}_0(z, \bar{z}) \right], \quad \text{such that } \alpha \left[\hat{\mathcal{Z}}^{h^*, \bar{h}^*}(z, \bar{z}) \right] = 1$$

$$\text{and } \alpha \left[\hat{\mathcal{Z}}^j(z, \bar{z}) \right] \geq 0, \quad \alpha \left[\hat{\mathcal{Z}}^{\bar{j}}(z, \bar{z}) \right] \geq 0 \quad \text{for } j \in \mathbb{Z},$$

$$\text{and } \alpha \left[\hat{\mathcal{Z}}^{h, \bar{h}}(z, \bar{z}) \right] \geq 0 \quad \text{for } (h, \bar{h}) \in \mathcal{P}$$

- This gives the maximum bound of the degeneracy of the state with (h^*, \bar{h}^*) .

$$d(h^*, \bar{h}^*) \leq -\alpha \left[\hat{\mathcal{Z}}_0(z, \bar{z}) \right]$$

- Extremal Functional Method [Paulos, El-Showk 14]

- Suppose the degeneracies of all primaries saturated the maximum bound. Then, the modular bootstrap equation is reduced to the below form.

$$\sum_{j=J_{min}}^{J_{max}} \left(d(j) \beta^* \left[\hat{\mathcal{Z}}^j(z, \bar{z}) \right] + \bar{d}(j) \beta^* \left[\hat{\mathcal{Z}}^{\bar{j}}(z, \bar{z}) \right] \right) + \sum_{h, \bar{h} \in \mathcal{P}} d(h, \bar{h}) \beta^* \left[\hat{\mathcal{Z}}^{h, \bar{h}}(z, \bar{z}) \right] = 0$$

- For the primaries, the above reduced equation forces :

$$d(h, \bar{h}) \beta^* \left[\hat{\mathcal{Z}}^{h, h}(z, \bar{z}) \right] = 0, \quad \text{for } \forall (h, \bar{h}) \in \mathcal{P}.$$


Idea : **Find the states such that $\beta^* \left[\hat{\mathcal{Z}}^{h, h}(z, \bar{z}) \right] = 0!$** (Otherwise, $d(h, \bar{h}) = 0$.)

- Spectrum Analysis

1. Apply the EFM and find the states such that make $\beta^* \left[\hat{\mathcal{Z}}^{h, h}(z, \bar{z}) \right] = 0$.
2. For those states, find the corresponding maximal degeneracies.
3. *Assuming every primaries hit the maximal degeneracies*, find the consistent modular invariant partition function.

- F_4 example

- The EFM analysis applied to the hypothetical CFT with $c = \frac{26}{5}$,



- From the EFM analysis, the data of spin-0 and spin-1 low-lying primaries are,

$$\Delta_{j=0} \in \left\{ \frac{6}{5} + 2n, 2 + 2n \mid n \in \mathbb{Z}_{\geq 0} \right\}, \quad \Delta_{j=1} \in \left\{ \frac{11}{5} + 2n, 3 + 2n \mid n \in \mathbb{Z}_{\geq 0} \right\}.$$

- The solutions to the second order MDE with $c = \frac{26}{5}$ gives :

$$f_0^{c=26/5}(q) = q^{-\frac{13}{60}} \left(1 + 52q + 377q^2 + 1976q^3 + 7852q^4 + \dots \right),$$

$$f_1^{c=26/5}(q) = q^{\frac{3}{5} - \frac{13}{60}} \left(26 + 299q + 1702q^2 + 7475q^3 + 27300q^4 + \dots \right).$$

- F_4 example (continued)

- For each low-lying primaries, the maximum degeneracies are,

(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg
$(\frac{3}{5}, \frac{3}{5})$	676.0000	(1, 1)	2704.0000	(1, 0)	52.00028
$(\frac{3}{5}, \frac{8}{5})$	7098.0001	(2, 1)	16848.001	(2, 0)	324.0007
$(\frac{3}{5}, \frac{13}{5})$	35802.002	(3, 1)	80444.061	(3, 0)	1547.0091
$(\frac{8}{5}, \frac{8}{5})$	74529.0001	(2, 2)	104976.005	(4, 0)	5499.0126

- The relation between *partition function* and *reduced partition function* is given by,

$$\hat{Z}_{F_4}(q, \bar{q}) = |\tau|^{\frac{1}{2}} \eta(\tau)^2 \bar{\eta}(\bar{\tau})^2 Z_{F_4}(q, \bar{q}) - \underbrace{(1-q)(1-\bar{q})}_{\text{Vaccum contribution}}$$

- The partition function of $(F_4)_1$ WZW model is known :

$$Z_{F_4}(q, \bar{q}) = |f_0^{c=26/5}(q)|^2 + |f_1^{c=26/5}(q)|^2$$

This perfectly agree with the numerical result.

- The Result Summary ($c \leq 8$)

- In case of $(G_2)_1$, $(F_4)_1$ and $(E_7)_1$ WZW model, its modular invariant partition function is known. In terms of the solutions to the second order MDE, they are written as [Gannon 92]

$$Z_{G_2}(q, \bar{q}) = |f_0^{c=14/5}(q)|^2 + |f_1^{c=14/5}(q)|^2$$

$$Z_{F_4}(q, \bar{q}) = |f_0^{c=26/5}(q)|^2 + |f_1^{c=26/5}(q)|^2$$

$$Z_{E_7}(q, \bar{q}) = |f_0^{c=7}(q)|^2 + |f_1^{c=7}(q)|^2$$

and in case of $(E_6)_1$ WZW model,

$$Z_{E_6}(q, \bar{q}) = f_0^{c=6}(q)\bar{f}_0^{c=6}(\bar{q}) + 2f_1^{c=6}(q)\bar{f}_1^{c=6}(\bar{q})$$

For them, we checked the spectral analysis successfully reproduce the known partition function.

- The $(A_1)_1$, $(A_2)_1$, $(G_2)_1$, $(D_4)_1$ and $(E_8)_1$ WZW models are realized by the scalar gap problem (Collier, Lin, Yin 16), it turns out that it also realized by the **twist gap problem**.

- $(E_{7,1/2})_1$ WZW model?

- $E_{7,1/2}$ is non-simple Lie algebra, its subalgebra is E_7 . It splits into $E_7 \oplus 56 \oplus \mathbb{R}$.
- The degeneracy analysis at $c = \frac{38}{5}$ gives, [Cohen, Man de 96], [Landsberg, Manivel 06]

(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg
$(\frac{4}{5}, \frac{4}{5})$	3249.0004	(1, 1)	36100.000	(1, 0)	190.00412
$(\frac{4}{5}, \frac{9}{5})$	59565.012	(2, 1)	501600.00	(2, 0)	2640.0481
$(\frac{9}{5}, \frac{9}{5})$	1092025.06	(2, 2)	6969600.01	(3, 0)	19285.021

- The solutions to the second order MDE with $c = 38/5$ are given by,

$$f_0^{c=38/5}(q) = q^{-\frac{19}{60}} \left(1 + 190q + 2831q^2 + 22306q^3 + 129276q^4 + \dots \right),$$

$$f_1^{c=38/5}(q) = q^{\frac{4}{5} - \frac{19}{60}} \left(57 + 1102q + 9367q^2 + 57362q^3 + 280459q^4 + \dots \right).$$

- If there is $(E_{7,1/2})_1$ WZW model, the modular invariant partition function may have the following diagonal form.


$$Z_{E_{7,1/2}}(q, \bar{q}) = |f_0^{c=\frac{38}{5}}(q)|^2 + |f_1^{c=\frac{38}{5}}(q)|^2$$

- Examine ECFTs via the modular bootstrap
 - CLAIM : Twist gap problem **realize the ECFTs with $c = 24, 48$** on the boundary.
 - The partition function of $c = 24$ ECFT is obtained by the solutions to **the third order MDE**, while the $c = 48$ partition function is realized by **the fourth order MDE**.

$$c = 24 : Z_{c=24}(q, \bar{q}) = J(q)\bar{J}(\bar{q})$$

$$c = 48 : Z_{c=48}(q, \bar{q}) = (J(q)^2 - 393767)(\bar{J}(\bar{q})^2 - 393767)$$

- The EFM analysis suggests that all of them have the states with integer Δ .



- We find that the results of the numerical analysis perfectly matched to the modular invariant partition functions.

- CFTs without Kac-Moody symmetry
 - In the mathematics, the corresponding vertex operator algebra was constructed.

Exceptional Vertex Operator Algebras and the Virasoro Algebra

Michael P. Tuite

$C = 8, d_2 = 155$: This can be realized as the fixed point free lattice VOA V_L^+ (fixed under the automorphism lifted from the reflection isometry of the lattice L) for the rank 8 even lattice $L = \sqrt{2}E_8$. The automorphism group is $O_{10}^+(2).2$ [G].

$C = 16, d_2 = 2295$: The VOA V_L^+ for the rank 16 Barnes-Wall even lattice $L = \Lambda_{16}$ whose automorphism group is $2^{16}.O_{10}^+(2)$ [S].

$C = 23\frac{1}{2}, d_2 = 96255$: This can be realized as the integrally graded subVOA of Höhn's Baby Monster Super VOA VB^{\natural} whose automorphism group is the Baby Monster group \mathbb{B} [Ho2].

- With an ansatz $f_0(q) = q^\alpha(a_0 + a_2q^2 + a_3q^3 + a_4q^4 + \dots)$, the solutions to the third order MDE with $c = 8, c = 16$ and $c = \frac{47}{2}$ are given by,

$$f_0^{c=8}(q) = q^{-1/3} \left(1 + 156q^2 + 1024q^3 + 6790q^4 + 32768q^5 + \dots \right)$$

$$f_0^{c=16}(q) = q^{-2/3} \left(1 + 2296q^2 + 65536q^3 + 1085468q^4 + \dots \right)$$

$$f_0^{c=47/2}(q) = q^{-47/48} \left(1 + 96256q^2 + 9646891q^3 + 366845011q^4 + \dots \right)$$

- The partition function of $c = 8$ CFT without Kac-Moody symmetry
- The degeneracy analysis without conserved current of $j = 1$ gives,

(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg
$(\frac{1}{2}, \frac{1}{2})$	496.0000000	(1, 1)	33728.00000	(2, 0)	155.000000
$(\frac{1}{2}, \frac{3}{2})$	17360.00000	(2, 1)	505920.0000	(3, 0)	868.000000
$(\frac{3}{2}, \frac{3}{2})$	607600.0009	(2, 2)	7612825.000	(4, 0)	5610.00000

- The other two solutions to the third order MDE with $c = 8$ are :

$$f_{h=1/2}(\tau) = a_0 q^{1/6} \left(1 + 36q + 394q^2 + 2776q^3 + 15155q^4 + \dots \right),$$

$$f_{h=1}(\tau) = a_1 q^{2/3} \left(1 + 16q + 136q^2 + 832q^3 + 4132q^4 + \dots \right)$$

- Our numerical results suggest that the modular invariant partition function reads,

$$\begin{aligned}
 Z_{c=8} &= f_{h=0}^{c=8}(\tau) \bar{f}_{h=0}^{c=8}(\bar{\tau}) + 496 f_{h=1/2}^{c=8}(\tau) \bar{f}_{h=1/2}^{c=8}(\bar{\tau})|_{a_0=1} + 33728 f_{h=1}^{c=8}(\tau) \bar{f}_{h=1}^{c=8}(\bar{\tau})|_{a_1=1}. \\
 &= 1 + \underbrace{496}_{1+155+340} q^{\frac{1}{2}} \bar{q}^{\frac{1}{2}} + \underbrace{17856}_{2 \times 155 + 2 \times 868 + 15810} q^{\frac{3}{2}} \bar{q}^{\frac{1}{2}} + \underbrace{33728}_{2108+31620} q \bar{q} + \underbrace{539648}_{539648} q^2 \bar{q} + \dots
 \end{aligned}$$

- The partition function of $c = 16$ CFT without Kac-Moody symmetry
- The degeneracy analysis without conserved current of $j = 1$ gives, gives,

(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg
$(\frac{3}{2}, \frac{3}{2})$	32505856.0032	(1, 1)	134912.0000	(2, 0)	2295.00000
$(\frac{3}{2}, \frac{5}{2})$	1657798656.0001	(2, 1)	18213120.00	(3, 0)	63240.0000
$(\frac{3}{2}, \frac{7}{2})$	34228666368.005	(2, 2)	2464038225.003	(4, 0)	1017636.00

- The other two solutions to the third order MDE with $c = 16$ are :

$$f_{h=1} = b_0 q^{1/3} \left(1 + 136q + 4132q^2 + 67712q^3 + 770442q^4 + \dots \right),$$

$$f_{h=3/2} = b_1 q^{5/6} \left(1 + 52q + 1106q^2 + 14808q^3 + 147239q^4 + \dots \right)$$

- Our numerical results suggest that the modular invariant partition function reads,

$$Z_{c=16} = f_{h=0}^{c=16}(\tau) \bar{f}_{h=0}^{c=16}(\bar{\tau}) + 134912 f_{h=1}^{c=16}(\tau) \bar{f}_{h=1}^{c=16}(\bar{\tau})|_{b_0=1} + 32505856 f_{h=3/2}^{c=16}(\tau) \bar{f}_{h=3/2}^{c=16}(\bar{\tau})|_{b_1=1}.$$

$$= 1 + \underbrace{2296}_{2 \times 1 + 186 + 2108} q^2 + \underbrace{65536}_{2 \times 1 + 186 + 14756 + 50592} q^3 + \underbrace{134912}_{186 + 340 + 868 + 22858 + 110670} q\bar{q} + \dots$$

- Baby Monster CFT [Höhn 07]

- The degeneracies with $c = \frac{47}{2}$ reads,

(h, \bar{h})	Max. Deg (h, \bar{h})	(h, \bar{h})	Max. Deg
$(\frac{3}{2}, \frac{3}{2})$	19105641.026984403127	$(\frac{5}{2}, \frac{5}{2})$	1298173112605.3499336
$(2, 2)$	9265025041.322733803	$(\frac{31}{16}, \frac{31}{16})$	9265217540.6086142750
$(\frac{5}{2}, \frac{3}{2})$	4980203754.2560961756	$(\frac{47}{16}, \frac{31}{16})$	1011288637613.8107313

- The three solutions to the third order MDE with $c = \frac{47}{2}$ are given by,

$$f_{h=3/2}^{c=47/2} = q^{25/48} a_1 \left(1 + \frac{785}{3} q + \frac{44393}{3} q^2 + 418441 q^3 + \frac{23301881}{3} q^4 + \dots \right)$$

$$f_{h=31/16}^{c=47/2} = q^{23/24} a_2 \left(1 + \frac{5177}{47} q + 4372 q^2 + 100627 q^3 + 1625207 q^4 + \dots \right)$$

- Corresponding modular invariant partition function reads,

$$Z_{c=47/2} = f_{h=0}^{c=47/2}(\tau) \bar{f}_{h=0}^{c=47/2}(\bar{\tau}) + f_{h=3/2}^{c=47/2}(\tau) \bar{f}_{h=3/2}^{c=47/2}(\bar{\tau}) \Big|_{a_1=\sqrt{4371}} + f_{h=31/16}^{c=47/2}(\tau) \bar{f}_{h=31/16}^{c=47/2}(\bar{\tau}) \Big|_{a_2=\sqrt{96256}}$$

$$= 1 + \underbrace{96256}_{1+96255} q^2 + \underbrace{9646891 q^3}_{2 \times 1 - 4371 + 2 \times 96255 + 9458750} + \underbrace{19105641 q^{3/2} \bar{q}^{-3/2}}_{1+96255+9458750+9550635} + \dots$$

- Comments on the “dual” CFT description
 - Ising model versus Babymonster CFT

	c	h_1	h_2
Ising model	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{16}$
Baby monster CFT	$\frac{47}{2}$	$\frac{3}{2}$	$\frac{31}{16}$
Sum	24	2	2

Ising model and Baby monster CFT are related via bilinear relation : [Hampapura, Mukhi 16]

$$j(\tau) - 744 = \chi_{\text{VB}_{(0)}^h}(\tau) \chi_{\text{vac}}^{\text{Ising}}(\tau) + \chi_{\text{VB}_{(1)}^h}(\tau) \chi_{h=\frac{1}{2}}^{\text{Ising}}(\tau) + \chi_{\text{VB}_{(3)}^h}(\tau) \chi_{h=\frac{1}{16}}^{\text{Ising}}(\tau)$$

- $c = 8$ and $c = 16$ CFTs without Kac-Moody symmetry are related by bilinear relation :

$$j(\tau) - 744 = \left(f_{h=0}^{c=8} \right) \left(f_{h=0}^{c=16} \right) + \left(f_{h=1}^{c=8} \Big|_{a_1=\sqrt{33728}} \right) \left(f_{h=1}^{c=16} \Big|_{b_0=\sqrt{134912}} \right) + \left(f_{h=1/2}^{c=8} \Big|_{a_0=\sqrt{496}} \right) \left(f_{h=3/2}^{c=16} \Big|_{b_1=\sqrt{32505856}} \right)$$

Application to the \mathcal{W} -algebra cases

- Bootstrapping with \mathcal{W} -algebra

- In case of the $\mathcal{W}(2,3)$ -algebra, we have spin-3 generator W_n . The corresponding fugacity $p = e^{2\pi iz}$ should be introduced in the character.

$$\begin{aligned}\chi_{(h,w;c)}(\tau, z) &= \text{Tr}_{h,w}(q^{L_0 - \frac{c}{24}} p^{W_0}) \\ &= \text{Tr}_{h,w}(q^{L_0 - \frac{c}{24}}) + \alpha_1 \text{Tr}_{h,w}(q^{L_0 - \frac{c}{24}} W_0) + \alpha_2 \text{Tr}_{h,w}(q^{L_0 - \frac{c}{24}} W_0^2) + \dots\end{aligned}$$

The modular transformation property is only known up to W_0^2 order. [Iles, Watts 14]

We will focus on the **unrefined character** which means W_0 -zeroth order character.


$$\chi_0(\tau) = \frac{q^{-\frac{c-2}{24}}(1-q)^3(1+q)}{\eta(\tau)^2}, \quad \chi(\tau) = \frac{q^{h-\frac{c-2}{24}}}{\eta(\tau)^2}$$

because of the null states $\langle 0|L_1 L_{-1}|0\rangle = 0$, $\langle 0|W_1 W_{-1}|0\rangle = 0$ and $\langle 0|W_2 W_{-2}|0\rangle = 0$.

- Assuming the non-vacuum module is non-degenerate, the unrefined character of rank- r $\mathcal{W}(d_1, d_2, \dots, d_r)$ -algebra is given by,



$$\chi_0(\tau) = \frac{q^{-\frac{c-N+1}{24}}}{\eta(\tau)^{N-1}} \prod_{j=1}^r \prod_{i=1}^{f_j-1} (1-q^i), \quad \chi(\tau) = \frac{q^{h-\frac{c-N+1}{24}}}{\eta(\tau)^{N-1}}.$$

- The Numerical Bounds(Twist Gap)



- The numerical bound at $c \geq r$ is identical to the one obtained from the Virasoro character. This results suggest that the unitary irreducible representations of $\mathcal{W}(d_1, d_2, \dots, d_r)$ -algebra do not contain any nontrivial null states when $c \geq r$.

- Numerical bound with Rank-3 \mathcal{W} -algebra



- $(c = 3, \Delta = \frac{3}{4})$ sits on the numerical boundary that obtained using the unrefined character of rank-3 $\mathcal{W}(2, 3, 4)$ algebra. Note that $c = 3$ is not in the list of the two-character RCFTs.
- The hypothetical CFT with $c = 3$ can be identified to the $(A_3)_1$ WZW model. This theory is realized by third order MDE, with Kac-Moody symmetry.
- CLAIM : The twist gap problem with $\mathcal{W}_{2,3,4}$ algebra EXCLUSIVELY realize $(A_3)_1$ WZW model on the numerical bound!

- Spectral Analysis on $(A_3)_1$ WZW model

- The maximal degeneracies are :

(h, \bar{h})	Max. Deg	(h, \bar{h})	Max. Deg
$(\frac{3}{8}, \frac{3}{8})$	32.00000	$(\frac{1}{2}, \frac{1}{2})$	36.000000
$(\frac{3}{8}, \frac{11}{8})$	96.00000	$(\frac{1}{2}, \frac{3}{2})$	48.00000
$(\frac{11}{8}, \frac{11}{8})$	288.01585	$(\frac{3}{2}, \frac{3}{2})$	64.11818

- The characters of (A_3) WZW model reads,

$$\chi_{[0]}^{A_3}(q) = q^{-\frac{1}{8}} \left(1 + 15q + 51q^2 + 172q^3 + 453q^4 + 1128q^5 + \dots \right),$$

$$\chi_{[1]}^{A_3}(q) = q^{\frac{3}{8} - \frac{1}{8}} \left(4 + 24q + 84q^2 + 248q^3 + 648q^4 + 1536q^5 + \dots \right),$$

$$\chi_{[2]}^{A_3}(q) = q^{\frac{1}{2} - \frac{1}{8}} \left(6 + 26q + 102q^2 + 276q^3 + 728q^4 + 1698q^5 + \dots \right),$$

- We find the maximal degeneracies are perfectly agree with the degeneracies in the below partition function.

$$Z_{A_3}(q, \bar{q}) = |\chi_{[0]}^{A_3}(q)|^2 + 2|\chi_{[1]}^{A_3}(q)|^2 + |\chi_{[2]}^{A_3}(q)|^2$$

• Conclusion and Outlook

- The **twist gap problem** with **holomorphic currents** ($j \geq 1$) successfully realize two-character RCFTs and three-character RCFTs on the numerical bound. The various RCFTs include level-one WZW models and extremal conformal field theories.
- When the holomorphic currents are included from $j = 2$, the CFTs without Kac-Moody symmetry are realized on the numerical boundary. It include $c = 8$, $c = 16$ CFTs and baby monster CFT. **We suggest the modular invariant partition function of those theories** based on the numerical results.

The coefficients in partition function can be decomposed by the dimension of irrep of the $O_{10}^+(2)$ or **baby monster group**. They are expected to be a underlying symmetry of the three special theories.

- The numerical analysis extended to the \mathcal{W} -algebra cases, using the unrefined character. The numerical bounds suggest the absence of the degenerate states in unitary irreducible representation when $c \geq r$.
- Application to the supersymmetric cases : Can we examine the super WZW models, super extremal conformal field theory? Unexpected super-RCFTs?