

Introduction to nonequilibrium physics

Jae Dong Noh

December 19, 2016

Preface

This is a note for the lecture given in the **2016 KIAS-SNU Physics Winter Camp** which is held at KIAS in December 17–23, 2016.

Most systems in nature are in a nonequilibrium state. In contrast to the thermal equilibrium systems, dynamics is important for nonequilibrium systems. The lecture covers the Langevin equation and the Fokker-Planck equation formalism for the study of dynamics of both equilibrium and nonequilibrium systems. It also covers the basic concepts of stochastic thermodynamics and the fluctuation theorems. This lecture is mainly based on the books [Gar10, Ris89, VK11].

Contents

1	Brownian motion	1
1.1	Einstein's theory	1
1.2	Langevin's theory	2
1.3	Perrin's experiment	3
2	Langevin equation	5
2.1	Brownian motion	6
2.2	Ornstein-Uhlenbeck process	7
2.3	Wiener process	7
2.4	Integral form of the Langevin equation	8
2.5	Numerical integration of the Langevin equation	9
3	Fokker-Planck Equation	10
3.1	Markov process and Chapman-Komogorov equation	10
3.2	Fokker-Planck equation	13
3.2.1	Random walks	15
3.2.2	Brownian motion	15
3.2.3	Three-Dimensional Brownian Motion in an External Force (Kramers problem)	16

Contents

3.2.4	Biased diffusion in a ring	17
3.3	Path integral or Onsager-Machlup formalism	18
3.4	Detailed balance and the Fluctuation Dissipation Theorem	19
4	Stochastic Thermodynamics	20
4.1	Stochastic energetics	21
4.1.1	Underdamped Langevin system	21
4.1.2	Overdamped Langevin system	22
4.1.3	Overdamped Langevin system with time-dependent potential	23
4.2	Nonequilibrium entropy	24
4.3	Time-irreversibility	25
4.4	Fluctuation theorem for the entropy production	27
4.5	FT for the work	28
4.6	More	28
5	Summary	29
	References	31
	Bibliography	31

1

Brownian motion

Brown in 1827 observed small pollen grains suspended in water at finite temperature T through a microscope. He observed an irregular motion of the small particles, which is later called the Brownian motion.

1.1 Einstein's theory

Einstein in 1905 explain the nature of the Brownian motion by assuming the stochastic nature of interactions between Brownian particles and fluid molecules. Let $f(x, t)dx$ be the fraction of Brownian particles between x and $x + dx$ at time t . If a Brownian particle is kicked by fluid molecules during the time interval τ , it is displaced by the amount of Δ . It is reasonable to assume that Δ is a random variable with a distribution function

1 Brownian motion

$\phi(\Delta) = \phi(-\Delta)$. Then, one obtains that

$$f(x, t + \tau) = \int_{-\infty}^{\infty} f(x - \Delta, t) \phi(\Delta) d\Delta \quad (1.1)$$

Expansion upto $O(\Delta^2)$ leads to the diffusion equation

$$\frac{\partial f}{\partial t} = D_d \frac{\partial^2 f}{\partial x^2} \quad (1.2)$$

with the diffusion constant $D_d = \frac{1}{2\tau} \int \Delta^2 \phi(\Delta) d\Delta$. Its solution is given by

$$f(x, t) = \frac{1}{\sqrt{4\pi D_d t}} e^{-(x-x_0)^2/(4D_d t)}. \quad (1.3)$$

The root mean square displacement is then scales as

$$\sqrt{\langle (x - x_0)^2 \rangle} = \sqrt{2D_d t} \propto t^{1/2}. \quad (1.4)$$

The $t^{1/2}$ dependence is the hallmark of the diffusive motion.

1.2 Langevin's theory

Langevin in 1906 suggested another explanation for the Brownian motion. He incorporated the effects of the random interaction into the Newton's equation of motion, and

proposed the following equation:

$$m \frac{d^2x}{dt^2} = -\gamma \frac{dx}{dt} + X(t). \quad (1.5)$$

This is the first appearance of the stochastic differential equation. In a rather crude way, Langevin derived the same relation in (1.4). He also derived the relation

$$D_d = \frac{k_B T}{\gamma} \quad (1.6)$$

known as the Einstein relation.

1.3 Perrin's experiment

Perrin in 1908 recorded the trajectory of a colloidal particle of radius $0.53\mu m$ every 30 seconds, and confirmed the diffusive motion of the Brownian particle experimentally. He was awarded the Nobel Prize in Physics in 1926.

1 *Brownian motion*




Figure 1.1: Perrin's experiment

2

Langevin equation

We are interested in a particle (or system of particles) in a thermal heat bath at temperature T . The thermal heat bath itself is a collection of many molecules and interact with the system in a complex way. At the phenomenological level, it is reasonable to assume that the heat bath provides a damping force on average and a fluctuating random force. The resulting Langevin equation is given by

$$v = \frac{dx}{dt} \quad , \quad m \frac{dv}{dt} = f(x, t) - \gamma v + \zeta(t), \quad (2.1)$$

where $\zeta(t)$ is the Gaussian distributed white noise satisfying

$$\langle \zeta(t) \rangle = 0 \quad , \quad \langle \zeta(t) \zeta(t') \rangle = 2D \delta(t - t'). \quad (2.2)$$

More rigorous derivation of the Langevin equation is found in the lecture of Prof. Yeo given in **PSI 2014** (<http://psi.kias.re.kr/2014/>).

2.1 Brownian motion

The Langevin equation for a free Brownian particle is given by

$$m\dot{v} = -\gamma v + \zeta(t) \text{ with the initial condition } v(0) = v_0. \quad (2.3)$$

It has the formal solution

$$v(t) = v_0 e^{-(\gamma/m)t} + \frac{1}{m} \int_0^t e^{-(\gamma/m)(t-t')} \zeta(t') dt'. \quad (2.4)$$

Using the formal solution, we can calculate the average value of various quantities. The average velocity decays exponentially as $\langle v(t) \rangle = v_0 e^{-(\gamma/m)t}$. The two-time correlation function is given by

$$\langle (v(t) - \langle v(t) \rangle)(v(t') - \langle v(t') \rangle) \rangle = \frac{D}{\gamma m} \left(e^{-(\gamma/m)|t-t'|} - e^{-(\gamma/m)(t+t')} \right). \quad (2.5)$$

The equal-time correlation function in the infinite time limit converges to

$$\langle v^2 \rangle = \frac{D}{\gamma m}. \quad (2.6)$$

The kinetic energy of a free particle in thermal equilibrium is given by $E = \frac{m}{2} \langle v^2 \rangle_{eq} = \frac{1}{2} k_B T$. By imposing the Einstein relation

$$D = \gamma k_B T \quad (2.7)$$

the Langevin equation describes the equilibrium dynamics.

2.2 Ornstein-Uhlenbeck process

Consider a harmonic oscillator in a heat bath at temperature T , whose Langevin equation is given by

$$\dot{x} = v \quad , \quad m\dot{v} = -kx - \gamma v + \zeta(t) . \quad (2.8)$$

When the damping coefficient is very large or the mass is very small, the system is in the overdamped regime. The Langevin equation then becomes

$$\gamma\dot{x} = -kx + \zeta(t) . \quad (2.9)$$

A stochastic process governed by the linear Langevin equation is called the OU process. In fact, the Brownian motion in the previous section is also an OU process for the velocity v . Using the connection, one can easily evaluate $\langle x^2 \rangle_{eq}$.

The general form of the OU process is given by

$$\dot{q}_i = \sum_j a_{ij} q_j + \zeta_i(t) \quad (2.10)$$

where $\langle \zeta_i(t) \rangle = 0$ and $\langle \zeta_i(t) \zeta_j(t') \rangle = 2D_{ij} \delta(t - t')$ with the noise matrix $D_{ij} = D_{ji}$.

2.3 Wiener process

The Wiener process, the simplest stochastic process, has the Langevin equation

$$\dot{W} = \eta(t) \quad \text{with} \quad \langle \eta(t) \rangle = 0 \quad \text{and} \quad \langle \eta(t) \eta(t') \rangle = \delta(t - t') \quad (2.11)$$

2 Langevin equation

The solution is given by

$$W(t) = W_0 + \int_0^t \eta(t') dt'. \quad (2.12)$$

One can show that

$$\langle W(t) \rangle = W_0 \quad \text{and} \quad \langle (W(t) - W_0)^2 \rangle = t. \quad (2.13)$$

The Wiener process describes the trajectory of a random walker.

Consider the displacement $dW(t) \equiv W(t + dt) - W(t) = \int_t^{t+dt} \zeta(t') dt'$ during the infinitesimal time interval dt . It is Gaussian distributed random variable with

$$\langle dW(t) \rangle = 0 \quad \text{and} \quad \langle dW(t)^2 \rangle = dt. \quad (2.14)$$

It indicates that $dW = O(\sqrt{dt})$.

2.4 Integral form of the Langevin equation

The Langevin equation of the form $\dot{q} = f(q, t) + \zeta(t)$ can be written as

$$dq = f(q, t)dt + dW(t) \quad (2.15)$$

where $dq \equiv q(t + dt) - q(t)$.

2.5 Numerical integration of the Langevin equation

The solution of the Langevin equation in (2.15) can be simulated numerically. For a stochastic trajectory in the time interval $[0 : t]$, first discretize the time as $t_{i=0, \dots, N} = i(\Delta t)$ with $\Delta t = t/N$. Then, $q_i = q(t_i)$ are found from the recursion relation

$$q_{i+1} = q_i + f(q_i, t_i)\Delta t + \sqrt{2D(\Delta t)} r_i, \quad (2.16)$$

where r_i is an independent random variable of zero mean ($\langle r_i \rangle = 0$) and unit variance ($\langle r_i^2 \rangle = 1$). This method is called the Euler method. This is a straightforward extension of the Euler method for the ordinary differential equation. Advanced numerical methods for the stochastic differential equation are found in Ref. [GSH88].

[Exercise] Integrate numerically the Langevin equation in (2.8) with $k = \gamma = k_B T = m = 1$ starting from the initial condition $x(0) = v(0) = 0$. Confirm the equipartition theorem $\langle x(\infty)^2 \rangle = \langle v(\infty)^2 \rangle = 1$.

3

Fokker-Planck Equation

Figure 3.1 displays 6 different realizations of the Wiener process trajectory. Due to the stochastic nature of the Langevin equation, $W(t)$ is a random variable fluctuating from sample to sample. Thus, it is meaningful to consider the probability distribution $P(W(t) = X|W(0) = 0)$. In this chapter, we study the Fokker-Planck equation that governs the time evolution of the probability distribution function.

3.1 Markov process and Chapman-Kolmogorov equation

Consider a time-dependent random variable $X(t)$ generated from a stochastic process. It is fully characterized by the joint probability densities $p(x_1, t_1; x_2, t_2; x_3, t_3; \dots)$. It describes how much probable it is to measure the values x_1, x_2, x_3, \dots at time $t_1 > t_2 > t_3 > \dots$. In terms of the joint probability densities, one can also define conditional probability

3 Fokker-Planck Equation




Figure 3.1: Trajectories of the Wiener process.

3 Fokker-Planck Equation

densities

$$p(x_1, t_1; x_2, t_2; \dots | y_1, \tau_1; y_2, \tau_2; \dots) = \frac{p(x_1, t_1; x_2, t_2; \dots; y_1, \tau_1; y_2, \tau_2; \dots)}{p(y_1, \tau_1; y_2, \tau_2; \dots)}, \quad (3.1)$$

where the times are ordered as $t_1 > t_2 > \dots > \tau_1 > \tau_2 > \dots$.

A stochastic process is called the **Markov process**, named after a Russian mathematician Andrey Markov (1956–1922), if the conditional probability is determined entirely by the knowledge of the most recent condition, i.e.,

$$p(x_1, t_1; x_2, t_2; \dots | y_1, \tau_1; y_2, \tau_2; \dots) = p(x_1, t_1; x_2, t_2; \dots | y_1, \tau_1). \quad (3.2)$$

In the Markov process, the future is determined by the present not by the past. A memory effect destroys the Markov property.

In the Markov process, any joint probability can be factorized as

$$p(1; 2; \dots; n) = p(1|2)p(2|3) \dots p(n-1|n)p(n). \quad (3.3)$$

Therefore, the transition probability density $p(x, t|y, t')$ fully characterizes a Markov process. Once you know the transition probability densities and the initial probability density, any joint probability density is determined.

The Markov property is a very strong condition. The transition probability for a Markov process should satisfy the Chapman-Kolmogorov equation

$$p(x_1, t_1|x_3, t_3) = \int dx_2 p(x_1, t_1|x_2, t_2)p(x_2, t_2|x_3, t_3). \quad (3.4)$$

It is named after a British mathematician Sydney Chapman (1888–1970) and a Russian mathematician Andrey Kolmogorov (1903–1987).

3.2 Fokker-Planck equation

A stochastic process governed by the Langevin equation

$$dx = A(x, t)dt + B dW(t) \quad (3.5)$$

belongs to the class of Markov processes. For convenience, we assume that B does not depend on x . The Langevin equation generates a stochastic time trajectory $x(t)$. Let $f(x)$ be an arbitrary well-behaved function. Then, the infinitesimal difference $df[x(t)] \equiv f[x(t + dt)] - f[x(t)]$ is given by

$$\begin{aligned} df[x(t)] &= f'[x(t)]dx(t) + \frac{1}{2}f''[x(t)]dx(t)^2 + \dots \\ &= f'[x(t)] \{A[x(t), t]dt + B dW(t)\} + \frac{1}{2}f''[x(t)]B^2 dW(t)^2 + \dots \end{aligned} \quad (3.6)$$

Taking the average, one obtains that

$$\frac{d}{dt} \langle f[x(t)] \rangle = \left\langle A[x(t), t]f'[x(t)] + \frac{1}{2}B^2 f''[x(t)] \right\rangle. \quad (3.7)$$

3 Fokker-Planck Equation

The averages are expressed in terms of the transition probability density $p(x, t|x_0, t_0)$ as

$$\begin{aligned} \int dx f(x) \partial_t p(x, t|x_0, t_0) &= \int dx \left[A(x, t) f'(x) + \frac{1}{2} b f''(x) \right] p(x, t|x_0, t_0) \\ &= \int dx f(x) \left[-\partial_x (A p) + \frac{1}{2} \partial_x^2 (B p) \right]. \end{aligned} \quad (3.8)$$

Since this relation is valid for any function $f(x)$, the transition probability should satisfy

$$\frac{\partial}{\partial t} p(x, t|x_0, t_0) = -\frac{\partial}{\partial x} [A(x, t) p(x, t|x_0, t_0)] + \frac{1}{2} \frac{\partial^2}{\partial x^2} [B^2 p(x, t|x_0, t_0)]. \quad (3.9)$$

This is called the Fokker-Planck equation, named after a Dutch physicist Adriaan Fokker (1887–1972) and a German physicist Max Planck (1858–1947). The first and second terms are called the *drift* and the *diffusion* terms, respectively. One can also derive the Fokker-Planck equation from the Kramers-Moyal expansion, which is not covered in this lecture.

When there are many variables $\mathbf{x} = (x_1, x_2, \dots)$ coupled through the Langevin equation

$$dx_i = A_i(\mathbf{x}, t) dt + \sum_j B_{ij} dW_j(t), \quad (3.10)$$

the Fokker-Planck equation for $p(\mathbf{x}, t|x_0, t_0) = p(\mathbf{x}, t)$ is given by

$$\partial_t p = -\sum_i \partial_i [A_i p] + \sum_{i,j} \partial_i \partial_j [D_{ij} p] = \mathcal{L}_{FP} p \quad (3.11)$$

3 Fokker-Planck Equation

with the noise matrix $D_{ij} = \frac{1}{2} \sum_k B_{ik} B_{jk}$ or $D = \frac{1}{2} B B^t$.

The Fokker-Planck equation can be written in the form of the continuity equation

$$\partial_t p + \sum_i \partial_i J_i = 0 \quad (3.12)$$

with the probability current

$$J_i = A_i p - \sum_j \partial_j (D_{ij} p) . \quad (3.13)$$

The probability current consists of the drift current and the diffusion current.

3.2.1 Random walks

The motion of a random walker is described by $dx = \sqrt{2D} dW$ with $A = 0$ and $B = \sqrt{2D}$. The corresponding Fokker-Planck equation is then given by

$$\frac{\partial p}{\partial t} = D \frac{\partial^2 p}{\partial x^2} . \quad (3.14)$$

This is the diffusion equation Einstein obtained.

3.2.2 Brownian motion

The Langevin equation for the Brownian motion is given by $m dv = -\gamma v dt + \sqrt{2\gamma T} dW$. Comparing it with (3.5), one finds that $A = -(\gamma/m)v$ and $B = \sqrt{2\gamma T/m^2}$. So, the

3 Fokker-Planck Equation

Fokker-Planck equation for $P(v, t)$ becomes

$$\frac{\partial}{\partial t} p(v, t) = \frac{\gamma}{m} \frac{\partial}{\partial v} [vp(v, t)] + \frac{\gamma T}{m^2} \frac{\partial^2}{\partial v^2} p(v, t) . \quad (3.15)$$

The time-dependent solution can be obtained by solving the partial differential equation using the Fourier transformation technique. Here, I only calculate the steady-state distribution function $p_{ss}(v) = \lim_{t \rightarrow \infty} p(v, t)$. The steady-state distribution should satisfy

$$J_{ss}(v) = -\frac{\gamma}{m} v p_{ss}(v) - \frac{\gamma T}{m^2} p'_{ss}(v) = J_0 (= \text{constant}) . \quad (3.16)$$

The steady-state current J_0 should be 0. Otherwise, the probability would build up at $v = \infty$ for positive J_0 or $v = -\infty$ for negative J_0 . The current-free condition yields that

$$p_{ss}(v) = \frac{1}{\sqrt{2\pi T/m}} e^{-\frac{mv^2}{2T}} . \quad (3.17)$$

This is the equilibrium Maxwell-Boltzmann distribution.

3.2.3 Three-Dimensional Brownian Motion in an External Force (Kramers problem)

The Langevin equations are

$$\begin{pmatrix} dx \\ dv \end{pmatrix} = \begin{pmatrix} v \\ -\frac{\gamma v}{m} + \frac{F}{m} \end{pmatrix} dt + \begin{pmatrix} 0 \\ \frac{2\gamma T}{m} d\mathbf{W} \end{pmatrix} \quad (3.18)$$

3 Fokker-Planck Equation

The corresponding Fokker-Planck equation for $p(\mathbf{x}, \mathbf{v}, t)$ is

$$\partial_t p = \left[-\nabla_{\mathbf{x}} \cdot \mathbf{v} + \nabla_{\mathbf{v}} \cdot \left(\frac{\gamma}{m} \mathbf{v} - \frac{\mathbf{F}}{m} \right) + \frac{\gamma T}{m^2} \nabla_{\mathbf{v}}^2 \right] p. \quad (3.19)$$

In general, even the steady state distribution is hard to find.

The steady state can be found in the special case with the conservative force

$$\mathbf{F}(\mathbf{x}) = -\nabla_{\mathbf{x}} V(\mathbf{x}) \quad (3.20)$$

with a scalar potential $V(\mathbf{x})$. The steady state distribution is given by the equilibrium Boltzmann distribution

$$p_{ss}(\mathbf{x}, \mathbf{v}) = \frac{1}{Z} \exp \left[-\frac{1}{T} \left(\frac{1}{2} m \mathbf{v}^2 + V(\mathbf{x}) \right) \right]. \quad (3.21)$$

3.2.4 Biased diffusion in a ring

Consider a Langevin equation $d\theta = f_0 + \sqrt{2T} dW$ for an angle variable $0 \leq \theta < 2\pi$. The Fokker-Planck equation for $p(\theta, t)$ is given by

$$\partial_t p = -\partial_{\theta} J_{\theta} \quad (3.22)$$

with the probability current $J_{\theta} = (f_0 - T\partial_{\theta})p(\theta, t)$. In the steady state, the current should be a constant J_0 independent of θ . Therefore, $Tp'_{ss}(\theta) = f_0 p_{ss}(\theta) - J_0$, which has a general solution $p_{ss}(\theta) = c e^{(f_0/T)\theta} + J_0/f_0$ with a constant c . The continuity $p_{ss}(0) = p_{ss}(2\pi)$ requires that $c = 0$ and the normalization $\int d\theta p_{ss}(\theta) = 1$ yields that $J_0 = \frac{f_0}{2\pi}$.

3.3 Path integral or Onsager-Machlup formalism

Consider the Langevin equation $dx = A(x)dt + B dW$ for a single variable x . Generalization to multivariate problems is straightforward. We want to calculate the transition probability for an infinitesimal time interval, $p(x', t + dt|x, t)$. A particle can move from x to $x' = x + (x' - x)$ only when the Langevin noise takes the right value $dW = (x' - x - A(x)dt)/B$. Recalling that dW is Gaussian distributed with mean zero and variance dt , we obtain that

$$p(x', t + dt|x, t) = \frac{1}{\sqrt{2\pi B^2 dt}} \exp\left(-\frac{[x' - x - A(x)dt]^2}{2B^2 dt}\right). \quad (3.23)$$

The transition probability over a finite time interval $p(x, t|x_0, t_0)$ is obtained by using the Markov property. We divide the time interval $[0 : t]$ into N sub-intervals of duration $\tau = t/N$. Then, the transition probability is written as

$$p(x, t|x_0, 0) = \left[\prod_{i=1}^{N-1} \int dx_i \right] \left[\prod_{i=0}^{N-1} p(x_{i+1}, t_{i+1}|x_i, t_i) \right]. \quad (3.24)$$

In the $N \rightarrow \infty$ limit, it becomes

$$\begin{aligned} p(x, t|x_0, 0) &= \lim_{N \rightarrow \infty} \left[\prod_{i=0}^{N-1} \int \frac{dx_i}{\sqrt{2\pi B^2 \tau}} \right] \exp\left(-\sum_{i=0}^{N-1} \frac{[x_{i+1} - x_i - A(x_i)\tau]^2}{2B^2 \tau}\right) \\ &= \int [\mathcal{D}x(s)] e^{-\int ds [\dot{x}(s) - A(x(s))]^2 / (2B^2)} \end{aligned} \quad (3.25)$$

3 Fokker-Planck Equation

The multivariate system with $dx_i = A_i dt + B_{ij} dW_j$ has the transition probability

$$\begin{aligned} p(\mathbf{x}, t | \mathbf{x}_0, 0) &= \int [\mathcal{D}\mathbf{x}(s)] e^{-\frac{1}{4} \int_0^t ds [\dot{\mathbf{x}}(s) - \mathbf{A}(\mathbf{x}(s))]_i D_{ij}^{-1} [\dot{\mathbf{x}}(s) - \mathbf{A}(\mathbf{x}(s))]_j} \\ &= \int [\mathcal{D}\mathbf{x}(s)] e^{-A[\mathbf{x}(s)]} \end{aligned} \quad (3.26)$$

with the action functional $A[\mathbf{x}(s)]$ and $D_{ij} = \frac{1}{2} \sum_k B_{ik} B_{jk}$.

The Onsager-Machlup [OM53] formalism allows us to write down the formal expression for the conditional path probability density

$$\Pi[\mathbf{x}[s] | \mathbf{x}_0] \propto e^{-A[\mathbf{x}(s)]} \quad (3.27)$$

3.4 Detailed balance and the Fluctuation Dissipation Theorem

A Fokker-Planck system with the steady state solution $p_{ss}(x) = e^{-\Phi(x)}$ is said to satisfy the detailed balance if

$$p(x', t' | x, t) p_{ss}(x) = p(\epsilon x, t | \epsilon x', t') p_{ss}(\epsilon x') \quad (3.28)$$

with the parity operator ϵ .

Please refer to the lecture note of Prof. Yeo of PSI 2014.

4

Stochastic Thermodynamics

We have studied the Langevin equation and the Fokker-Planck equation formalism for general stochastic systems. We are ready to study thermodynamics of such systems at the level of microscopic stochastic paths. We will learn how to define thermodynamic quantities such as heat, work, and entropy, and investigate some symmetry properties of thermodynamic systems. References for this chapter are Refs. [Sek10, Sei05, Kwo15].




Figure 4.1: Thermal system

4.1 Stochastic energetics

4.1.1 Underdamped Langevin system

Consider a particle (or a system of particles) in thermal contact with a heat bath of temperature T . The Langevin equation is given by

$$\begin{aligned} x &= v dt \\ m dv &= f_c(x) dt + f_{nc}(x) dt - \gamma v dt + \sqrt{2\gamma T} dW(t) , \end{aligned} \quad (4.1)$$

where $f_c(x) = -\frac{dV(x)}{dx}$ is the conservative force, and f_{nc} is a time-independent non-conservative force. The internal energy of the system is defined as

$$E = \frac{1}{2} m v^2 + V(x) . \quad (4.2)$$

It is a fluctuating variable since v is a stochastic variable. For the change of the kinetic energy during the infinitesimal time step dt , we need to calculate

$$dv^2 \equiv (v + dv)^2 - v^2 = 2v dv + (dv)^2 = 2 \frac{v + (v + dv)}{2} dv = 2v \circ dv \quad (4.3)$$

Note that one should not neglect $(dv)^2$ because dv involves dW whose square is of the order of dt . We can use the form of normal calculus by introducing the special multiplication scheme denoted as \circ . This is called the Stratonovich calculus: $X(t) \circ dY(t) = \frac{X(t) + X(t+dt)}{2} dY(t)$.

4 Stochastic Thermodynamics

The energy change is then given by

$$dE = mv \circ dv - f_c(x) \circ dx . \quad (4.4)$$

Eliminating $f_c = mdv/dt - f_{nc} + \gamma vdt - \sqrt{2\gamma T}dW/dt$ using the Langevin equation, one obtains

$$\begin{aligned} dE &= mv \circ dv - v \circ \left[mdv - f_{nc}dt - (-\gamma vdt + \sqrt{2\gamma T}dW) \right] \\ &= f_{nc} \circ dx + v \circ (-\gamma vdt + \sqrt{2\gamma T}dW) \\ &= dW + dQ. \end{aligned} \quad (4.5)$$

This is the first law of thermodynamics for the Langevin system with the work and the heat

$$\begin{aligned} dW &= f_{nc} \circ dx \\ dQ &= v \circ \left(-\gamma vdt + \sqrt{2\gamma T} dW \right) = F_B \circ dx. \end{aligned} \quad (4.6)$$

The expression for the heat looks quite intuitive. It is the “work” done by the force F_B from the heat bath [Sek10].

4.1.2 Overdamped Langevin system

Langevin equation is given by

$$\gamma dx = f_c dt + f_{nc} dt + \sqrt{2\gamma T} dW = f_{tot} dt + \sqrt{2\gamma T} dW . \quad (4.7)$$

4 Stochastic Thermodynamics

The first law of thermodynamics becomes as

$$dV = dW + dQ \quad (4.8)$$

where

$$\begin{aligned} dW &= f_{nc} \circ dx \\ dQ &= F_B \circ dx = -(f_c + f_{nc}) \circ dx . \end{aligned} \quad (4.9)$$

4.1.3 Overdamped Langevin system with time-dependent potential

Suppose that the potential energy includes a time dependence parameter

$$V = V(x, \lambda(t)) . \quad (4.10)$$

The parameter is called the protocol. You may think of a harmonic oscillator with a time dependent spring constant. The first law of thermodynamics becomes

$$dV = dW_{nc} + dW_J + dQ \quad (4.11)$$

where

$$dW_J = \frac{\partial V}{\partial \lambda} \frac{d\lambda}{dt} dt \quad (4.12)$$

is called the Jarzynski work.

4.2 Nonequilibrium entropy

What is the entropy of the nonequilibrium system? We adopt the Shannon's information entropy [Sha48] as the thermodynamic entropy. It is defined as

$$\langle S_{sys}(t) \rangle = - \int dx p(q, t) \ln p(q, t) . \quad (4.13)$$

It is the ensemble average of the instant entropy

$$S_{sys}(q(t), t) = - \ln p(q(t), t) \quad (4.14)$$

of a system being at $q(t)$ at time t .

When a system evolve in time following a stochastic path $[q(s)]$ for $t_i \leq s \leq t_f$, the system entropy changes as

$$\Delta S_{sys}[q] = - \ln \frac{p(q(t_f), t_f)}{p(q(t_i), t_i)} . \quad (4.15)$$

The entropy change of the heat bath is given by the Clausius form

$$\Delta S_B[q] = \frac{-\Delta Q}{T} . \quad (4.16)$$

The total entropy change is then given by $\Delta S_{tot} = \Delta S_{sys} + \Delta S_B$. Does it satisfy the second law of thermodynamics?

4.3 Time-irreversibility

We will show that the entropy is directed related to the **time-irreversibility**. Specifically, I explain the concept in the context of the overdamped system with the Langevin equation in (4.7). Using the Onsager-Machlup formalism, we can write down the formal expression for the probability density that the system would follow a path $[x(s)]$ connecting x at time t to $x' = x + dx$ at time $t + dt$. It is given by

$$\text{Prob}[x(s)] = p_i(x)\Pi[x(s)|x_i] \propto p(x, t) \exp \left[-\frac{\gamma dt}{4T} \left\{ \frac{(x' - x)}{dt} - \frac{f_{tot}(x)}{\gamma} \right\}^2 \right], \quad (4.17)$$

where $f(x) = f_c(x) + f_{nc}(x)$ is the total force.

Imagine the time-reversed path $[x^R(s)]$ connecting x' at time t to x at time $t + dt$. The probability density of the reverse path is given by

$$\text{Prob}[x^R(s)] \propto p(x', t + dt) \exp \left[-\frac{\gamma dt}{4T} \left\{ \frac{(x - x')}{dt} - \frac{f_{tot}(x')}{\gamma} \right\}^2 \right]. \quad (4.18)$$

The irreversibility is measured with the log ratio of the path probability densities:

$$\begin{aligned} \ln \frac{\text{Prob}[x]}{\text{Prob}[x^R]} &= \ln \frac{p(x, t)}{p(x', t + dt)} + \frac{dt}{4T} \left[2 \frac{(x' - x)}{dt} (f(x) + f(x')) - \frac{f(x)^2 - f(x')^2}{\gamma} \right] \\ &= dS_{sys} + \frac{f \circ dx}{T} = dS_{sys} + \frac{-dQ}{T} = dS_{sys} + dS_B \end{aligned} \quad (4.19)$$




Figure 4.2: Forward and reverse paths

That is to say, the total entropy change is the same as the irreversibility. Note that we have degrees of freedom in choosing the initial and final probability distributions.

4.4 Fluctuation theorem for the entropy production

It is straightforward to show that

$$\langle e^{-\Delta S} \rangle = \int dS P(S) e^{-S} = 1 \quad (4.20)$$

for any Langevin system. This is called the integral fluctuation theorem (IFT). Applying the Jensen's inequality $\langle e^{-x} \rangle \geq e^{-\langle x \rangle}$, one can derive the second law of thermodynamics

$$\langle \Delta S \rangle \geq 0. \quad (4.21)$$

When the system is in the steady state, we have more powerful fluctuation theorem for the probability distribution $P(S)$.

$$\frac{P(S)}{P(-S)} = e^S. \quad (4.22)$$

This is called the detailed fluctuation theorem (DFT). The DFT holds only when the system satisfies the involution property.

4.5 FT for the work

Consider a system with a time-dependent potential energy $V(x, \lambda(t))$. If it is in the thermal equilibrium state initially, then we have the IFT

$$\langle e^{-W/T} \rangle = e^{-\Delta F/T} . \quad (4.23)$$

This is called the Jarzynski equality [Jar97]. We also have the DFT

$$\frac{P_F(W)}{P_R(-W)} = e^{-\Delta(W-F/T)} . \quad (4.24)$$

which is called the Crooks fluctuation theorem [Cro99].

4.6 More

Maxwell's demon, heat engine, ...

5

Summary

- Langevin equation : stochastic processes, stochastic differential equation, numerical integration
- Fokker-Planck equation : Markov process, Chapman-Kolmogorov equation, Ito and Stratonovich, Onsager-Machlup path integral formalism
- Stochastic thermodynamics : stochastic energetics, irreversibility, entropy production, fluctuation theorem, ...

5 Summary

[Project] The motion of a rotary motor is described by the overdamped Langevin equation

$$\frac{d\theta}{dt} = -\frac{dV(\theta)}{d\theta} + f_0 + \eta(t),$$

where $0 \leq \theta < 2\pi$ is the angular position of the motor, $V(\theta) = \cos\theta$ is the periodic potential, and f_0 is the constant applied torque.

(1) Write down the corresponding Fokker-Planck equation for the probability distribution function $P(\theta, t)$ and solve it to find the stationary state distribution $P_{ss}(\theta)$. Compare it with the equilibrium Boltzmann distribution.

(2) Find the average angular velocity $\Omega = \lim_{t \rightarrow \infty} \frac{1}{t} \langle (\theta(t) - \theta(0)) \rangle$ as a function of the applied torque f_0 .

(3) Prepare your system in the equilibrium state with f_0 and then turn on the torque at time $t = 0$. Solve the Langevin equation numerically upto time $t = \tau$ and measure the work done by the external torque

$$W(\tau) = \int_0^\tau f_0 \dot{\theta}(t') dt' .$$

By performing the numerical simulations many times, you can construct the probability distribution function for the work $P(W)$. Test the integral and detailed fluctuation theorems by computing $\langle e^{-W} \rangle$ and $P(W)/P(-W)$.

Bibliography

- [Cro99] GE Crooks. Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Phys. Rev. E*, 60(3):2721, 1999.
- [Gar10] Crispin Gardiner. *Stochastic Methods*. A Handbook for the Natural and Social Sciences. Springer, New York, fourth edition, 2010.
- [GSH88] A Greiner, W Strittmatter, and J Honerkamp. Numerical integration of stochastic differential equations. *J. Stat. Phys.*, 51(1-2):95–108, April 1988.
- [Jar97] C Jarzynski. Nonequilibrium equality for free energy differences. *Phys. Rev. Lett.*, 78(14):2690–2693, 1997.
- [Kwo15] Chulan Kwon. Path-dependent entropy production. *J. Korean Phys. Soc.*, 67(5):785–791, September 2015.
- [OM53] L Onsager and S Machlup. Fluctuations and irreversible processes. *Phys. Rev.*, 91(6):1505–1512, 1953.
- [Ris89] Hannes Risken. *The Fokker-Planck Equation*. Methods of Solution and Applications. Springer-Verlag, Berlin, 1989.
- [Sei05] Udo Seifert. Entropy Production along a Stochastic Trajectory and an Integral Fluctuation Theorem. *Phys. Rev. Lett.*, 95(4):040602, July 2005.

Bibliography

- [Sek10] Ken Sekimoto. *Stochastic energetics*, volume 799 of *Lecture Notes in Physics*, Berlin Springer Verlag. 2010.
- [Sha48] C E Shannon. A Mathematical Theory of Communication. *The Bell System Technical Journal*, XXXII(4):378–423, July 1948.
- [VK11] N G Van Kampen. *Stochastic Processes in Physics and Chemistry*. Elsevier, August 2011.