

Probing origins of neutrino masses and baryon asymmetry

Takehiko Asaka (Niigata Univ.)

In collaboration with
Shintaro Eijima (EPFL, Switzerland)
Atsushi Watanabe (MPIK, Germany)

References: JHEP1303 (2013) 125
arXiv:1308.3550 (to appear in PTEP)

Introduction

- Neutrino mass scales
 - ▣ Atmospheric: $\Delta m_{\text{atm}}^2 \simeq 2.4 \times 10^{-3} \text{ eV}^2$
 - ▣ Solar : $\Delta m_{\text{sol}}^2 \simeq 7.5 \times 10^{-5} \text{ eV}^2$
- ⇒ Need for physics beyond the SM !
- Important questions:
 - ▣ ***“What is the origin of neutrino masses?”***
 - ▣ ***“How do we test it experimentally?”***

Extension by RH neutrinos ν_R

$$\delta L = i \overline{\nu}_R \partial_\mu \gamma^\mu \nu_R - F \overline{L} \nu_R \Phi - \frac{M_M}{2} \overline{\nu}_R \nu_R^c + \text{h.c.}$$

Minkowski '77

Yanagida '79

Gell-Mann, Ramond, Slansky '79

Glashow '79

- Seesaw mechanism ($M_D = F\langle\Phi\rangle \ll M_M$)

$$-L = \frac{1}{2} (\overline{\nu}_L, \overline{\nu}_R^c) \begin{pmatrix} 0 & M_D \\ M_D^T & M_M \end{pmatrix} \begin{pmatrix} \nu_L^c \\ \nu_R \end{pmatrix} + \text{h.c.} = \frac{1}{2} (\overline{\nu}, \overline{N}^c) \begin{pmatrix} M_\nu & 0 \\ 0 & M_M \end{pmatrix} \begin{pmatrix} \nu^c \\ N \end{pmatrix} + \text{h.c.}$$

$$M_\nu = -M_D^T \frac{1}{M_M} M_D$$

$$U^T M_\nu U = \text{diag}(m_1, m_2, m_3)$$

□ Light, active neutrinos ν

→ explain neutrino oscillations

□ Heavy neutrinos N ($N \simeq \nu_R$)

- Mass M_M
- Mixing $\Theta = M_D / M_M$

mixing in CC current $\nu_L = U \nu + \Theta N$




Where is
the scale
of mass?

Scale of Majorana mass

- The simplest case: one pair of ν_L and ν_R


$$M_\nu = -M_D^T \frac{1}{M_M} M_D \Rightarrow F^2 = M_M M_\nu / \langle \Phi \rangle^2$$



Scale of Majorana mass

- The simplest case: one pair of ν_L and ν_R


$$M_\nu = -M_D^T \frac{1}{M_M} M_D \Rightarrow F^2 = M_M M_\nu / \langle \Phi \rangle^2$$



Scale of Majorana mass

- The simplest case: one pair of ν_L and ν_R

$$M_\nu = -M_D^T \frac{1}{M_M} M_D \Rightarrow F^2 = M_M M_\nu / \langle \Phi \rangle^2$$



In this talk

Consider the minimal case with two RH neutrinos

- Lighter than charged kaon $M_{2,3} < m_K$
→ Test by Kaon decays ($K^+ \rightarrow \ell^+ N_I$) is possible
- Current status of (RH) heavy neutrinos
 - Region of successful baryogenesis
 - Constraints from direct search and cosmology
- Implication to $0\nu 2\beta$ decay
- Search for (RH) heavy neutrinos at T2K

When adding one more DM RH neutrino N_1 ,
the results can be applied to the ν MSM !!

TA, Blanchet, Shaposhnikov ('05),
TA, Shaposhnikov ('05)

§

Current status of heavy neutrinos

Baryogenesis via neutrino osc.

Oscillation of heavy neutrinos can be a source of BAU


Akhmedov, Rubakov, Smirnov ('98) / TA, Shaposhnikov ('05)

Shaposhnikov ('08), Canetti, Shaposhnikov ('10)

TA, Ishida ('10), Canetti, Drewes, Shaposhnikov ('12), TA, Eijima, Ishida ('12)

Canetti, Drewes, Shaposhnikov ('12), Canetti, Drewes, Frossard, Shaposhnikov ('12)

- CPV in oscillation and production generates asymmetries
- Asymmetries are separated into LH and RH leptons
- Asymmetry in LH leptons is converted into BAU



Yield of BAU depends on
Yukawa couplings $F_{\alpha I}$ and masses

Especially, CP violating parameters
and mass difference


$$T_{osc} \sim (M_0 M_N \Delta M)^{1/3}$$

Baryogenesis via neutrino osc.

10

Region accounting for $\frac{n_B}{s} = (8.55-9.00) \times 10^{-11}$

Canetti, Shaposhnikov '10




Baryogenesis via neutrino osc.

Region accounting for $\frac{n_B}{s} = (8.55-9.00) \times 10^{-11}$

- (1) quasi-degenerate
- (2) masses are

$M_N > 2.1 \text{ MeV (NH)}$
 $M_N > 0.7 \text{ MeV (IH)}$




Baryogenesis via neutrino osc.

Region accounting for $\frac{n_B}{s} = (8.55-9.00) \times 10^{-11}$

- (1) quasi-degenerate
- (2) masses are

$M_N > 2.1 \text{ MeV (NH)}$
 $M_N > 0.7 \text{ MeV (IH)}$


Such light RH neutrinos can be directly tested by experiments!



Direct search experiment


■ PS191 [Bernardi et al '86, '88]

Beam dump experiment
performed at CERN in 1984



- Production $\pi^+, K^+ \rightarrow e^+ N$
- Detection $N \rightarrow \ell^+ \ell^- \nu, \ell^- \pi^+$

■ Upper bounds mixing elements Θ \rightarrow Lower bound on lifetime of N



BBN constraint on lifetime



- Long-lived $N_{2,3}$ may spoil the success of BBN
 - ▣ Speed up the expansion of the universe
 - $\rho_{\text{tot}} = \rho_{\text{MSM}} + \rho_{N_{2,3}} \Rightarrow H^2 = \frac{\rho_{\text{tot}}}{3 M_P^2}$
 - p-n conv. decouples earlier \Rightarrow overproduction of ${}^4\text{He}$
 $n + \nu \leftrightarrow p + e^- , \dots$
 - ▣ Distortion of spectrum of active neutrinos
 - $N_{2,3} \rightarrow \nu \bar{\nu} \nu, e^+ e^- \nu, \dots$
 - Additional neutrinos may not be thermalized
- ⇒ Upper bound on lifetime

- Dolgov, Hansen, Rafflet, Semikoz ('00)
 - ▣ One family case: $\tau_N < 0.1 \text{ sec}$ for $M_N > m_\pi$

Constraints on light RH neutrinos

15

TA, Eijima '13



$M_N > 163$ MeV

$M_N = 188 - 269$ MeV
 $M_N > 285$ MeV



 §

Implication to $0\nu2\beta$ decay

Constraints on light RH neutrinos

17

TA, Eijima '13




$M_N > 163$ MeV

$M_N = 188 - 269$ MeV
 $M_N > 285$ MeV

Mixing elements in IH case

Mixing elements of heavy neutrinos $\Theta_{\alpha I}$



$$\Theta_{\alpha I} = \frac{\langle \Phi \rangle F_{\alpha I}}{M_I}$$

■ Mixing elements strongly depend on “ $\xi \sin \eta$ ”


$$|\Theta_e|^2 \simeq 1.20 \times 10^{-8} \left(\frac{\text{MeV}}{M_N} \right) (1.000 - 0.925\xi \sin \eta) X_\omega^2,$$

$$|\Theta_\mu|^2 \simeq 0.76 \times 10^{-8} \left(\frac{\text{MeV}}{M_N} \right) (1.000 + 0.895\xi \sin \eta - 0.250\xi \cos \eta \sin \delta + 0.092\xi \sin \eta \cos \delta) X_\omega^2,$$

$$|\Theta_\tau|^2 \simeq 0.50 \times 10^{-8} \left(\frac{\text{MeV}}{M_N} \right) (1.000 + 0.860\xi \sin \eta + 0.380\xi \cos \eta \sin \delta - 0.140\xi \sin \eta \cos \delta) X_\omega^2.$$

We find allowed range of Majorana phase !

Majorana phase in IH case




Majorana phase is restricted for $M_N < 350$ MeV!

$0\nu2\beta$ decays in IH

Effective neutrino mass from light and heavy neutrinos

$$m_{\text{eff}} = m_i U_{ei}^2 + f_\beta(M_I) M_I \Theta_{eI}^2 = [1 - f_\beta(M_N)] m_{\text{eff}}^\nu \quad \text{TA, Eijima, Ishida ('11)}$$

$$m_{\text{eff}}^\nu = \cos^2 \theta_{13} (m_1^2 \cos^4 \theta_{12} + m_2^2 \sin^4 \theta_{12} + 2m_1 m_2 \cos^2 \theta_{12} \sin^2 \theta_{12} \cos 2\eta)^{1/2}$$



- Heavy neutrinos give negative contribution to m_{eff}
- Constraint on η restricts the predicted range of m_{eff}

§



Search for heavy neutrinos at T2K

TA, Eijima, Watanabe
[JHEP1303 (2013) 125]

Constraints on light RH neutrinos

22

TA, Eijima '13




$M_N > 163$ MeV

$M_N = 188 - 269$ MeV
 $M_N > 285$ MeV

Search for heavy neutrinos at T2K

23



Search for heavy neutrinos at T2K

24


Production of N

$$K^+ \rightarrow \ell^+ + N \quad (\ell^- = e^-, \mu^-)$$


Detection of N

$$N \rightarrow \ell^- + \pi^+$$

- Estimate flux of N at ND280
- Count # of signal decay inside ND280
- Derive upper bounds on mixing angles



Sensitivity: PS191 vs T2K



T2K at 10^{21} POT has a better sensitivity than PS191
 $(0.86 \times 10^{19}$ POT) !

Signal vs Background

- Signal events: $N \rightarrow \ell^- + \pi^+$
- BG events: $\nu_\mu + n \rightarrow \mu^- + \pi^+ + n$ (CC- $n\pi^+$)
 $\nu_\mu + {}^{16}O \rightarrow \mu^- + \pi^+ + {}^{16}O$ (CC-coherent π^+)

To reduce BG,

- ▣ Use the invariant mass distribution of ℓ^- and π^+ since it has a peak at M_N for signal decay
- ▣ Use the low density part of detector filled with argon gas ($9m^3$) out of $61.25m^3$

See also the recent proposal to search for heavy neutrinos at the CERN SPS. (Shaposhnikov's talk)

arXiv:1310.1762

Summary

- We have considered the model with two right-handed neutrinos which are lighter than charged Kaon.

- Neutrino masses by seesaw mechanism
 - Baryogenesis via neutrino oscillations
 - Search in Kaon decays

- We have found the possible region for neutrino oscillations and BAU, allowed from search and cosmological constraints.

- Majorana phase is restricted in IH
→ Distinctive feature in $0\nu 2\beta$ decay

- We have discussed search for such right-handed neutrinos at near detector ND280 of T2K experiment

- Signal: $N \rightarrow e^- + \pi^+$, $\mu^- + \pi^+$ inside ND280
 - T2K at 10^{21} POT has a better sensitivity than PS191

Backup

Comparison

	PS191 [55, 56]	T2K [61]	MINOS [62]	MiniBooNE [63]	SciBooNE [64]
POT	0.86×10^{19}	10^{21}	10^{21}	10^{21}	10^{21}
(Distance) $^{-2}$	$(128\text{ m})^{-2}$	$(280\text{ m})^{-2}$	$(1\text{ km})^{-2}$	$(541\text{ m})^{-2}$	$(100\text{ m})^{-2}$
Volume	216 m^3	88 m^3	303 m^3	524 m^3	15.3 m^3
Events	1	9.9	2.7	15.8	13.5

Table 1. A comparison between PS191 and recent accelerator experiments. The item “Distance” means the distance between the beam target and the detector for each experiment. The item “Events” shows $\text{POT} \times (\text{Distance})^{-2} \times \text{Volume}$ in units of PS191. The POTs for the oscillation experiments are assumed to achieve 10^{21} .

T2K

2013/4/12: 6.39×10^{20} POT
 2013/5/8: 6.63×10^{20} POT

GOAL: 7.8×10^{21} POT

Neutrino Yukawa couplings for $N_{2,3}$

$$F = U_{\text{PMNS}} D_\nu^{1/2} \Omega D_N^{1/2} / \langle \Phi \rangle \quad (\text{in NH})$$

[Casas, Ibarra '01]

■ Parameters of active neutrinos

$D_\nu^{1/2} = \text{diag}(\sqrt{m_1} = 0, \sqrt{m_2}, \sqrt{m_3})$: active ν masses

$$U_{\text{PMNS}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}c_{12}s_{13}e^{i\delta} & c_{23}c_{12} - s_{23}s_{12}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}c_{12}s_{13}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{12}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}$$

$$\begin{pmatrix} 1 & & \\ & e^{i\eta} & \\ & & 1 \end{pmatrix}$$

■ Parameters of sterile neutrinos

$D_N^{1/2} = \text{diag}(\sqrt{M_2}, \sqrt{M_3})$: sterile ν masses

$$\Omega = \begin{pmatrix} 0 & 0 \\ \cos \omega & -\sin \omega \\ \xi \sin \omega & \xi \cos \omega \end{pmatrix}$$


ω : complex number
 $\xi = \pm 1$

Im ω

Effective neutrino mass


- $$m_{\text{eff}} = \sum_{i=1,2,3} m_i U_{ei}^2 + \sum_{I=1,2,3} f_\beta(M_I) M_I \Theta_{eI}^2$$

active neutrinos
sterile neutrinos



Takehiko Asaka (Niigata Univ.)

[Blennow, Fernandez-Martinez, Pavon, Mnendez '10]





15/11/2013

m_{eff} in the ν MSM

32

[TA, Eijima, Ishida '11]



- m_{eff} in the ν MSM is smaller than active ν 's one
- No significant constraint on Θ_{eI} in the ν MSM !